

FIG. 30J



FIG. 30K



FIG. 30L



FIG. 30M


Insect cell or fungi expressed interferon alpha-14C.
a-d, f, h, j-n, s, u, v-y = 0;
e, g, i, r, t (independently selected) = 0 or 1; z = 1.
1. endoglycanase
2. Galactosyltransferase,
UDP-Gal-linker-SA-CMP
3. ST3Gal3, transferrin
i (independently selected) = 0 or 1;
a-h, j-m, r-z = 0;
n = 1; R' = -Gal-linker-transferrin.

FIG. 30N





a-c, e, f (independently selected) = 0 or 1;
d, g = 0; R = polymer, glycoconjugate.

FIG. 30O



FIG. 30P



FIG. 30Q



FIG. 30R



FIG. 30S



FIG. 30T



FIG. 30U



FIG. 30V



FIG. 30W





a-c, e (independently selected) = 0 or 1;
d = 0; R = polymer, glycoconjugate.

FIG. 30X



FIG. 30Y



FIG. 30Z


E. coli expressed interferon alpha-mucin fusion protein.
a-e = 0.
1. GalNAc Transferase, UDP-GalNAc
2. CMP-SA-PEG, sialyltransferase
c, d, e (independently selected) = 0 or 1;
a, b = 0; R = PEG.

FIG. 30AA











a-c, e (independently selected) = 0 or 1;
d = 0; R = polymer, linker.

FIG. 30BB



FIG. 30CC



FIG. 30DD


E. coli expressed Interferon alpha (no fusion).
a-e, n = 0.
1. NHS-CO-linker-SA-CMP
2. ST3Gal3, transferrin
a-e = 0; n = 1; R' = linker-transferrin.

FIG. 30EE



a-d, i, r-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 or 1.
n, v-y = 0; z = 0 or 1; R = polymer

FIG. 31A

CHO, BHK, 293 cells, Vero expressed IF-beta
h = 1 to 3;
a-g, j-m, i (independently selected) = 0 or 1;
r-u (independently selected) = 0 or 1;
n, v-y = 0; z = 1.

1. Sialidase
2. CMP-SA-PEG, ST3Gal3

h = 1 to 3;
a-g, i (independently selected) = 0 or 1;
r-u (independently selected) = 0 or 1;
j-m, v-y (independently selected) = 0 or 1;
z = 1; n = 0; R = PEG.

FIG. 31B

Insect cell expressed IF-beta
a-d, f, h, j-n, s, u, v-y = 0;
e, g, i, r, t (independently selected) = 0 or 1;
z = 1.

1. GNT's 1&2, UDP-GlcNAc
2. Galactosyltransferase, UDP-Gal
2. CMP-SA-PEG, ST3Gal3,
buffer, salt

b, d, f, h, k, m, n, s, u, w, y = 0;
a, c, e, g, i, r, t (independently selected) = 0 or 1;
j, l, v, x (independently selected) = 0 or 1;
z = 1; R = PEG.

FIG. 31C



FIG. 31D



FIG. 31E



FIG. 31F



FIG. 31G



FIG. 31H



FIG. 31I



FIG. 31J



FIG. 31K

NSO expressed Ifn-beta.
a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; n, v-y = 0;
Sia (independently selected) = Sia or Gal.

↓
1. Sialidase and α-galactosidase
2. α-Galactosyltransferase, UDP-Gal
3. CMP-SA-PEG, ST3Gal3

a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; R = PEG
n = 0; v-y (independently selected) = 1,
when j-m (independently selected) is 1;

FIG. 31L

CHO, BHK, 293 cells, Vero expressed Ifn-beta.
a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; n, v-y = 0.

↓
1. Sialidase
2. CMP-SA-PEG (16 mol eq), ST3Gal3
3. CMP-SA, ST3Gal3

a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; n = 0;
v-y (independently selected) = 0 or 1; R = PEG.

FIG. 31M



FIG. 31N



FIG. 31O





a-d, i, p-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = modifying group;
R' = H, glycosyl group, modifying group, glycoconjugate.

FIG. 31P



FIG. 31Q



FIG. 31R


CHO, BHK, 293 cells, Vero expressed Ifn-beta.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA-linker-SA-CMP, ST3Gal3
2. ST3Gal3, desialylated transferrin.
3. CMP-SA, ST3Gal3
a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0;
v-y (independently selected) = 0 or 1;
R = linker-transferrin.

FIG. 31S





B ← {Glc-(Xyl)n}o

C ← {-Fuc}p

a-d, i, q-u (independently selected) = 0 or 1.
o, p (independently selected) = 0 or 1.
e-h, n (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 20.
v-y = 0;
R = modifying group, mannose, oligo-mannose, Sia-Lewis X, Sia-Lewis A..

FIG. 32A

BHK expressed Factor VII or VIIa
a-d, e, i, g, q, j, l, o, p (independently selected) = 0 or 1;
r, t = 1; f, h, k, m, s, u, v-y = 0; n = 0-4.

↓ 1. Sialidase
2. CMP-SA-PEG (16 mole eq), ST3Gal3

a-d, e, g, i, q, j, l, o, p (independently selected) = 0 or 1;
r, t = 1; f, h, k, m, s, u, w, y = 0; n = 0-4;
v, x, (independently selected) = 1,
when j, l (respectively, independently selected) is 1;
R = PEG.

FIG. 32B

CHO, BHK, 293 cells, Vero expressed Factor VII or VIIa
a-d, e, i, g, q, j, l, o, p (independently selected) = 0 or 1;
r, t = 1; f, h, k, m, s, u, v-y = 0; n = 0-4.

↓ 1. Sialidase
2. CMP-SA-PEG (1.2 mole eq), ST3Gal3
3. CMP-SA (8 mol eq), ST3Gal3

a-d, e, g, i, q, j, l, o, p (independently selected) = 0 or 1;
r, t = 1; f, h, k, m, s, u, w, y = 0; n = 0-4;
v or x, (independently selected) = 1,
when j or l, (respectively, independently selected) is 1;
R = PEG.

FIG. 32C


NSO expressed Factor VII or VIIa
a--u (independently selected) = 0 or 1;
v-y = 0; n = 0-4;
Sia (independently selected) = Sia or Gal.
1. Sialidase and α-galactosidase
2. Galactosyltransferase, UDP-Gal
3. CMP-SA-PEG, ST3Gal3
a-m, o-u (independently selected) = 0 or 1;
n = 0-4; v-y (independently selected) = 1,
when j-m (independently selected) is 1;
Sia = Sia; R = PEG.

FIG. 32D











a-d, i, n-u (independently selected) = 0 or 1.
bb, cc, dd, ee, ff, gg (independently selected) = 0 or 1.
e-h, aa (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 20.
v-z = 0; R = modifying group, mannose, oligo-mannose.

FIG. 33A

CHO, BHK, 293 cells, Vero expressed Factor IX
a-d, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, j-m, i, n, o, p, r-u (independently selected) = 0 or 1;
v-z, gg = 0.

1. Sialidase
2. CMP-SA-PEG, ST3Gal3

a-d, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, i, n, r-u (independently selected) = 0 or 1;
o, p, z = 0;
j-m, ee, v-y, gg (independently selected) = 0 or 1;
R = PEG.

FIG. 33B

CHO, BHK, 293 cells, Vero expressed Factor IX
a-d, n, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, j-m, i, o, p, r-u (independently selected) = 0 or 1;
v-z, gg = 0.

1. Sialidase
2. CMP-SA-PEG, ST3Gal3
3. ST3Gal1, CMP-SA

a-d, n, p, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, i, r-u (independently selected) = 0 or 1;
j-m, ee, v-y, gg (independently selected) = 0 or 1;
o, z = 0; R = PEG.

FIG. 33C

CHO, BHK, 293 cells, Vero expressed Factor IX
a-d, n, q, bb, cc, dd, ff = 1; e-h, aa = 1 to 4; ee, j-m, i, o, p, r-u (independently selected) = 0 or 1; v-z, gg = 0.

1. sialidase
2. Galactosyltransferase, UDP-Gal
3. CMP-SA, ST3Gal3
4. CMP-SA-PEG, ST3Gal1

a-d, n, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, i, r-u (independently selected) = 0 or 1; R = PEG;
o, v-y, gg = 0;
j-m, p, ee (independently selected) = 0 or 1, but when p = 1, z = 1.

FIG. 33D

CHO, BHK, 293 cells, Vero expressed Factor IX
a-d, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, j-m, i, n, o, p, r-u (independently selected) = 0 or 1;
v-z, gg = 0.

CMP-SA-PEG, ST3Gal3

a-d, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, i, n, r-u (independently selected) = 0 or 1; R = PEG;
o, p, z = 0; j-m, ee, v-y, gg (independently selected) = 0 or 1.

FIG. 33E

CHO, BHK, 293 cells, Vero expressed Factor IX
a-d, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, j-m, i, n, o, p, r-u (independently selected) = 0 or 1;
v-z, gg = 0.

1. CMP-SA-levulinate, ST3Gal3, buffer, salt
2. H_4N_2-PEG

a-d, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, i, n, r-u (independently selected) = 0 or 1;
o, p, z = 0; R = PEG;
j-m, ee, v-y, gg (independently selected) = 0 or 1.

FIG. 33F

CHO, BHK, 293 cells, Vero expressed Factor IX
a-d, n, q, bb, cc, dd, ff = 1;
e-h, aa = 1 to 4;
ee, j-m, i, o, p, r-u (independently selected) = 0 or 1;
v-z, gg = 0.

1. CMP-SA-PEG, α2,8-ST

a-d, q = 1; e-h = 1 to 4;
aa, bb, cc, dd, ee, ff, i, n, r-u (independently selected) = 0 or 1;
o, p, z = 0; R= PEG;
j-m, ee (independently selected) = 0 to 2;
v-y, gg (independently selected) = 1, when j-m (independently selected) is 2;

FIG. 33G





a-d, i, q-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0;
R = modifying group, mannose, oligo-mannose.

FIG. 34A



FIG. 34B



FIG. 34C



FIG. 34D



FIG. 34E



FIG. 34F



FIG. 34G



FIG. 34H



FIG. 34I


CHO, BHK, 293 cells, Vero expressed FSH.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA-linker-SA-CMP, ST3Gal3
2. ST3Gal1, desialylated chorionic gonadrophin (CG) produced in CHO.
3. CMP-SA, ST3Gal3, ST3Gal1
a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0;
v-y (independently selected) = 0 or 1;
R = linker-CG.

FIG. 34J





a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
v-z = 0;
R = polymer.

FIG. 35A

CHO, BHK, 293 cells, Vero expressed EPO
a-g, n, q = 1; h = 1 to 3;
j-m, i, o, p (independently selected) = 0 or 1;
r-u (independently selected) = 0 to1; v-z = 0

1. Sialidase
2. CMP-SA-PEG, ST3Gal3

a-g, n, q = 1; h = 1 to 3;
i, o, p (independently selected) = 0 or 1;
r-u (independently selected) = 0 or 1;
j-m, v-y (independently selected) = 0 or 1;
R = PEG; z = 0.

FIG. 35B

Insect cell expressed EPO
a-d, f, h, j-q, s, u, v-z = 0;
e, g, i, r, t (independently selected) = 0 or 1.

1. GNT's 1&2, UDP-GlcNAc
2. Galactosyltransferase, UDP-Gal
2. CMP-SA-PEG, ST3Gal3

b, d, f, h, k, m-q, s, u, w, y, z = 0;
a, c, e, g, i, r, t (independently selected)= 0 or 1;
j, l, v, x (independently selected) = 0 or 1;
R = PEG.

FIG. 35C

CHO, BHK, 293 cells, Vero expressed EPO
a-q, r-u (independently selected) = 0 or 1;
v-z = 0.

1. sialidase
2. Galactosyltransferase, UDP-Gal
3. CMP-SA, ST3Gal3
4. CMP-SA-PEG, ST3Gal1

a-h, n, q = 1;
i-m, o, r-u (independently selected) = 0 or 1;
v-y = 0; p, z = 0 or 1; R = PEG.

FIG. 35D

CHO, BHK, 293 cells, Vero expressed EPO
a-g, n, q = 1; h = 1 to 3;
j-m, i, o, p (independently selected) = 0 or 1;
r-u (independently selected) = 0 or 1;
v-z = 0

1. CMP-SA-PEG, ST3Gal3

a-g, n, q = 1; h = 1 to 3;
i, o, p (independently selected) = 0 or 1;
r-u (independently selected) = 0 to 1;
j-m, v-y (independently selected) = 0 or 1;
R = PEG; z = 0.

FIG. 35E



FIG. 35F



FIG. 35G



FIG. 35H



FIG. 35I



FIG. 35J



FIG. 35K



FIG. 35L



FIG. 35M



FIG. 35N



FIG. 35O



FIG. 35P



FIG. 35Q



FIG. 35R



FIG. 35S





a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
v-z = 0; aa, bb = 1; cc = 0;
R = polymer; R'' and R' = sugar-polymer or Fuc.

FIG. 35T



FIG. 35U



FIG. 35V


Insect cell expressed EPO
a-d, f, h, j-m, n-q, s, u, v-z = 0;
e, g, i, r, t (independently selected) = 0 or 1;
aa = 1; R" = Fuc.
1. mannosidases
2. galactosyltransferase, UDP-Gal-PEG
cc, e, i, r, v (independently selected) = 0 or 1;
bb, a-h, j-q, s-u, w-z = 0; aa = 1; R' = Gal-PEG.

FIG. 35W





a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
v-z = 0;
R = polymer.

FIG. 35X



FIG. 35Y



FIG. 35Z


CHO, BHK, 293 cells, Vero expressed NESP
a-g, n, q = 1; h = 1 to 3;
j-m, i, o, p (independently selected) = 0 or 1;
r-u (independently selected) = 0 to1; v-z = 0
1 CMP-SA, poly-α2,8-ST
a-g, n, q = 1; h = 1 to 3;
i, j-m, o, p, r-u, (independently selected) = 0 or 1;
v-z (independently selected) = 0-40; R = Sia.

FIG. 35AA







a-d, i, n-u, aa (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = polymer, glycoconjugate.

FIG. 36A



FIG. 36B



FIG. 36C



FIG. 36D



FIG. 36E



FIG. 36F



FIG. 36G







a-d, i, n-u, aa, bb, cc (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = modifying group, mannose, oligo-mannose.
R'= H, glycosyl residue, modifying group. glycoconjugate.

FIG. 36H

Insect cell expressed GM-CSF.
a-d, f, h, j-m, o, p, s, u, v-z = 0;
e, g, i, n, q, r, t, aa (independently selected) = 0 or 1.

↓ 1. GNT's 1,2,4,5, UDP-GlcNAc
2. Galactosyltransferase, UDP-Gal-PEG

a-i, n, q-u (independently selected) = 0 or 1;
j-m = 0; v-y (independently selected) = 1,
when e-h (independently selected) is 1;
R = PEG.

FIG. 36I

Yeast expressed GM-CSF.
a-p, z, cc = 0;
q-y, aa (independently selected) = 0 to 1;
bb = 1; R (branched or linear) = Man, oligomannose;
GalNAc = Man.

↓ 1. Endoglycanase
2. mannosidase (if aa = 1).
3. Galactosyltransferase, UDP-Gal-PEG

a-p, r-z, aa, bb = 0;
q, cc (independently selected) = 0 or 1;
R' = -Gal-PEG.

FIG. 36J


CHO, BHK, 293 cells, Vero expressed GM-CSF.
a--m, o-u, aa, bb (independently selected) = 0 or 1;
n, v-z, cc = 0.
1. sialidase
2. CMP-SA, ST3Gal3
2. CMP-SA-linker-SA-CMP, ST3Gal1
3. ST3Gal3, transferrin
a--m, p-u, z, aa (independently selected) = 0 or 1;
o, v-y, cc = 0; bb, n = 1; R = transferrin.

FIG. 36K





a-d, i, q-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = polymer.

FIG. 37A



FIG. 37B



FIG. 37C

NSO expressed Interferon gamma.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0;
Sia (independently selected) = Sia or Gal.

↓
1. Sialidase and α-galactosidase
2. α-Galactosyltransferase, UDP-Gal
3. CMP-SA-PEG, ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y (independently selected) = 1,
when j-m (independently selected) is 1;
R = PEG.

FIG. 37D

CHO, BHK, 293 cells, Vero expressed
Interferon gamma.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

↓
1. Sialidase
2. CMP-SA-PEG (16 mol eq),
ST3Gal3
3. CMP-SA, ST3Gal3

a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 37E



FIG. 37F



FIG. 37G





a-d, i, n, p-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0;
R = modifying group, mannose, oligo-mannose;
R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 37H

Insect or fungi cell expressed IF-gamma.
a-d, f, h, j-m, s, u, v-y = 0;
e, g, i, q, r, t (independently selected) = 0 or 1.

↓ 1. GNT's 1,2,4,5, UDP-GlcNAc
2. Galactosyltransferase, UDP-Gal-PEG

a-i, q-u (independently selected) = 0 or 1;
j-m = 0; v-y (independently selected) = 1,
when e-h (independently selected) is 1;
R = PEG.

FIG. 37I

Yeast expressed IF-gamma.
a-m = 0; q-y (independently selected) = 0 to 1; p = 1;
R (branched or linear) = Man, oligomannose.

↓ 1. Endoglycanase
2. Galactosyltransferase, UDP-Gal
3. CMP-SA-PEG, ST3Gal3

a-m, p-y = 0; n (independently selected) = 0 or 1;
R' = -Gal-Sia-PEG.

FIG. 37J



FIG. 37K



FIG. 37L

Insect or fungi cell expressed IF-gamma.
a-d, f, h, j-n, s, u, v-y = 0;
e, g, i, q, r, t (independently selected) = 0 or 1.

↓ 1. GNT's 1 & 2, UDP-GlcNAc-PEG

a-d, f, h, j-n, s, u, w, y = 0;
e, g, i, r, t, q (independently selected) = 0 or 1;
p = 1; v, x (independently selected) = 1,
when e, g (independently selected) is 1;
R = PEG.

FIG. 37M

CHO, BHK, 293 cells, Vero expressed
Interferon gamma.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.

↓ 1 . CMP-SA-PEG, α2,8-ST

a-d, i, q-u (independently selected) = 0 or 1;
e-h = 1; j-m (independently selected) = 0-2;
v-y (independently selected) = 1,
when j-m (independently selected) = 2;
R = PEG.

FIG. 37N





a-d, i, q-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = polymer.

FIG. 38A



FIG. 38B



FIG. 38C



FIG. 38D



FIG. 38E


CHO, BHK, 293 cells, Vero expressed α_1-antitrypsin.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA, α2,8-ST
a-d, i, q-u (independently selected) = 0 or 1; e-h = 1;
j-m (independently selected) = 0-20;
v-y (independently selected) = 0.

FIG. 38F





a-d, i, n, p-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0;
R = modifying group, mannose, oligo-mannose;
R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 38G



FIG. 38H



FIG. 38I


CHO, BHK, 293 cells, Vero expressed α_1-antitrypsin.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA-linker-Gal-UDP, ST3Gal3
2. Galactosyltransferase, transferrin treated with endoglycanase
a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0;
v-y (independently selected) = 0 or 1;
R = linker-transferrin.

FIG. 38J





a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
R = polymer;
R', R" (independently selected) = sugar, glycoconjugate.

FIG. 38K

Yeast expressed alpha-1 antitrypsin.
a-h, i-m, p, q = 0;
R (independently selected) = mannose, oligomannose, polymannose;
r-u, v-y (independently selected) = 0 or 1; n, o = 1.

↓ 1. endoglycanase
2. Galactosyltransferase, UDP-Gal-PEG

a-h, i-o, q, r-u, v-y = 0; p = 1.
R" = Gal-PEG.

FIG. 38L

Plant expressed alpha-1 antitrypsin.
a-d, f, h, j- m, s, u , v-y = 0;
e, g, i, q, r, t (independently selected) = 0 or 1;
n =1; R' = xylose

↓ 1. hexosaminidase,
2. alpha mannosidase and xylosidase
3. GlcNAc transferase, UDP-GlcNAc-PEG

a-d, f, h, j-n, s, u , v-y = 0;
e, g, i, r, t (independently selected) = 0;
q = 1; R' = GlcNAc-PEG.

FIG. 38M


CHO, BHK, 293 cells, Vero, transgenic animal expressed α_1 antitrypsin.
a-h, i-o, r-u (independently selected) = 0 or 1;
p, q, v-y = 0.
1. CMP-SA-PEG, ST3Gal3
a-h, i-o, r-u (independently selected) = 0 or 1;
p, q = 0; v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 38N